

Rapid Spectral Variability during the 2003 Outburst of V4641 Sgr (= SAX J1819.3-2525)¹

Dipankar Maitra^a, Charles Bailyn^a

^a*Yale University, New Haven, CT 06511, USA*

Abstract

The black hole candidate V4641 Sgr (= SAX J1819.3-2525) went through a brief outburst during 2003 Aug 01 to Aug 08. During the outburst, activity was noted in optical, radio as well as X-rays. Here we present results of Spectral and Temporal analysis of a pointed *Rossi X-ray Timing Explorer* (RXTE) observation of the source during the outburst. During this pointing we observed flaring activities with associated X-ray luminosity variations over factors of 5 or more in timescales of few tens of seconds. The observed flares during are intrinsically different in their spectral and temporal properties. We see evidence of variable column density during one of the flares. The spectral and temporal analyses of the data suggest occasional outflow/mass ejection.

Key words: accretion, accretion disks — stars, black holes — X-rays, binaries — individual (V4641 Sgr)

PACS: 01.30.Cc, 95.85.Nv, 97.80.Jp, 97.60.Lf, 98.62.Mw, 97.10.Gz, 04.70.-s

The compact binary system V4641 Sgr (= SAX J1819.3-2525) has been known to exhibit short, irregular outbursts, with highly variable lightcurve profile. The rapid time variability [1,2] in the lightcurve as well as observation of moving radio structure [3] during an outburst in 1999 led to its classification as a microblazar [4].

During early Aug 2003, signs of enhanced activity were reported by the VS-NET² group and shortly thereafter by the *Small and Moderate Aperture Research Telescope System* consortium telescopes at Cerro Tololo Inter-American Observatory (CTIO) in Chile [5]. In Fig. 1 we show the V band lightcurve of

Email address: maitra@astro.yale.edu (Dipankar Maitra).

¹ This work was supported by the National Science Foundation grant AST 0407063.

² <http://ooruri.kusastro.kyoto-u.ac.jp/mailman/listinfo/vsnet-alert>

V4641 Sgr during this outburst, as observed by the SMARTS 1.3m telescope. We used the CTIO observations to trigger the series of target-of-opportunity (ToO) X-ray observations. During four pointed RXTE observations we saw activity from the source. The observed X-ray spectrum during all these 4 pointings were hard in nature, showing large contribution of hard X-rays. Strong Fe $K\alpha$ fluorescent emission line near 6.5 keV was detected with large equivalent widths in the range of 700 - 1000eV.




Fig. 1. Optical V band lightcurve showing the 2003 outburst of V4641 Sgr. Four vertical lines on the top labelled 1-4 are the times when significant X-ray activity was observed by RXTE. The horizontal dotted line represents the mean quiescent brightness of 13.7 magnitude. The data were taken by the CTIO 1.3m telescope operated by the SMARTS Consortium.

In this article we concentrate, in particular, on second dwell where we observed two flaring activities with flare timescales of tens of seconds. In panel (a), (b), (c) and (d) of Fig. 2 we show the evolution of the source lightcurve, hardness ratio, color-magnitude diagram and color-color plane respectively during dwell (2). The distinctively opposite characters of the two flares are evident. While the first flare at ~ 250 seconds is essentially hard, the second flare during ~ 450 second is predominantly soft.

Since the Iron line dominates the low energy spectrum, estimating the continuum becomes difficult. We modelled the X-ray spectrum in the 10-25 keV region using the *pexrav* model [6] which calculates an exponentially cut off power law spectrum reflected from neutral material. The spectrum during the first 14-94 seconds is well fit by this model and a warm absorber with its column density (n_H) set to the standard value of 2.3×10^{21} atoms/cm $^{-2}$ for this source [7]. The fit model, when extended to softer energies, fails to account for the Fe $K\alpha$ line near 6.5 keV, but matches the continuum at the softest bands as shown in the left panel of Fig. 3. However, during the hard flare, the fits to high energy regimes of the spectrum (> 10 keV) largely overestimates the counts in softer energies as shown in the right panel of Fig. 3. In fact, we were unable to find reasonable fits with any physically motivated models with n_H fixed to its standard value. Allowing n_H results in much better fits. In Fig. 4 we show the variation of the fit parameter n_H during the hard flare.




Fig. 2. RXTE/PCA observations of V4641 starting MJD 52857.37 are shown. The data during the hard flare are shown by ■, those during soft flare are shown by ● and the rest of the dwell are shown by + symbols. Panel (a): The 3-20 keV lightcurve showing the rapid variability of this source that sets it apart from typical X-ray binaries. Panel (b): Variation of hardness-ratio with time. Hard band is 10.3-20.4 keV, medium band is 5.3-10.3 keV, soft band is 2.0-5.3 keV. Panel (c): Color-Luminosity plot showing the different regions occupied by the soft and hard flares. Panel (d): Color-color plot, also shows the while the soft flare occupies the most soft spectral state (lower-left corner), the hard flare essentially occupies the extreme hard state in the upper-right corner of the color-color diagram.




Fig. 3. Variation of n_H during the hard flare in dwell (2). Panel (a): The energy spectrum during 14-94 seconds. Panel (b): The spectrum during 222-254 seconds when the source was going through a hard flare. The solid histogram in both panels is a fit to the 10.0-25.0 keV spectrum with $n_H = 2.3 \times 10^{21} \text{ atoms/cm}^2$. Note that the extrapolated fit matches the continuum at lowest energies for panel (a) whereas it grossly overestimates the counts during the flare as shown in panel (b).

Conclusions

- V4641 Sgr went through a short lived, hard flaring outburst between 2003 Aug 01 through Aug 08, during which X-ray fluctuations by a factor of 10 on timescales of tens of seconds were seen.




Fig. 4. Variation of the fit parameter n_H during the hard flare. The dashed horizontal line represents the standard adopted value of n_H for this source.

- Compton reflection features like strong Fe $K\alpha$ line near 6.5 keV and curvature in spectrum at high energies were present in all observations.
- Spectral nature of some flares separated in time only by few minutes were intrinsically different.
- The thick warm envelope/outflow scenario [1,8] can explain the variation of column density during the second pointing although other scenarios like variation of partial covering fractions cannot be ruled out.

References

- [1] Revnivtsev, M., Gilfanov, M., Churazov, E., & Sunyaev, R. 2002, A&A, 391, 1013
- [2] Wijnands, R. & van der Klis, M. 2000, ApJ, 528, L93
- [3] Hjellming, R. M., et al. 2000, ApJ, 544, 977
- [4] Mirabel, I. F., & Rodriguez, L. F. 1994, Nature, 371, 46
- [5] Buxton, M., Maitra, D., Bailyn, C., Jeanty, L., & Gonzalez, D. 2003, The Astronomer's Telegram, 170, 1
- [6] Magdziarz, P., & Zdziarski, A. A. 1995, MNRAS, 273, 837
- [7] Dickey, J. M., & Lockman, F. J. 1990, ARA&A, 28, 215
- [8] Titarchuk, L., & Shrader, C. 2005, ApJ, 623, 362